



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

COROLLARY. Let  $\alpha = \frac{1}{2}\pi$ ; then  $\Delta = \frac{b^2}{2\pi}$ , the same as problem 26.

[NOTE.—By mistake in numbering the problems in this department, number 28 was omitted. The above problem and solution are inserted that problems be numbered consecutively. EDITOR.]

29. Proposed by JOHN DOLMAN, Jr., Philadelphia, Pennsylvania.

Neglecting perturbations, what is the average distance of the earth from the sun ?

Solution by WILLIAM HOOVER, A. M., Ph. D., Professor of Mathematics and Astronomy, Ohio University, Athens, Ohio.

The focus being the pole, the polar equation to the ellipse is

$$r = \frac{a(1-e^2)}{1-e\cos\theta} \dots \dots \dots (1).$$

I. The radii vectores being drawn at equal angular intervals,

$$m' = \frac{\int r d\theta}{\int d\theta} = a(1-e^2) \frac{\int_0^\pi \frac{d\theta}{1-e\cos\theta}}{\int_0^\pi d\theta} = a\sqrt{1-e^2} = b.$$

II. If  $x$  be the abscissa of any point on the curve, the focal distance is

$$r = a - ex \dots \dots \dots (2),$$

$$\text{and } m'' = \frac{\int_{-a}^{+a} (a-ex) dx}{\int_{-a}^{+a} dx} = a,$$

the points on the curve being so taken that their abscissas increase uniformly.

III. If the number of radii vectores depends upon the length of the curve,

$$m''' = \frac{\int r ds}{\int ds},$$

$ds$  being an element of the curve.

Also solved as I. above by Profs. F. P. MATZ, and O. W. ANTHONY, and as III. by Prof. G. B. M. ZERR.

## PROBLEMS.

37. Proposed by HENRY HEATON, M. Sc., Atlantic, Iowa.

Required the average area of all triangles two of whose sides are  $a$  and  $b$ .

38. Proposed by B. F. FINKEL, A. M., Professor of Mathematics and Physics, Drury College, Springfield, Missouri.

Two arrows are sticking in a circular target : show that the chance that their distance is greater than the radius of the target is  $3\sqrt{3}/4\pi$ . [From *Todhunter's Integral Calculus*, page 335.]

## MISCELLANEOUS.

Conducted by J. M. COLAW, Monterey, Va. All contributions to this department should be sent to him.

## SOLUTIONS OF PROBLEMS.

33. Proposed by Prof. ALEXANDER ROSS, C. E., Sebastopol, California,

From a point  $P$  without a square field  $ABCD$ , the distances  $PA$ ,  $PB$ , and  $PC$  measured to the corners are, respectively, 70, 40, and 60 chains. What is the area of the field ?

I. Solution by A. H. BELL, Hillsboro, Illinois, and A. H. HOLMES, Brunswick, Maine.

Let  $a > b > c$  equal the distances 70, 60, and 40, and let  $x = a$  side of the square field. Then  $\cos A = \frac{a^2 + x^2 - c^2}{2ax}$ , and this multiplied

by  $a = AF = \frac{a^2 + x^2 - c^2}{2x}$ .  $AF - AB = BF = EP = \frac{a^2 - c^2 - x^2}{2x}$  ;

then, also,  $BE = \frac{b^2 - c^2 - x^2}{2x}$ .


$$\therefore \frac{(a^2 - c^2 - x^2)^2 + (b^2 - c^2 - x^2)^2}{4x^2} = c^2 \dots\dots\dots(1).$$

$$x^4 - (a^2 + b^2)x^2 = c^2(a^2 + b^2) - \frac{a^4 + b^4 + 2c^4}{2} \dots\dots\dots(2).$$

$$\text{Area of square} = x^2 = \frac{1}{2} \left[ a^2 + b^2 \pm \sqrt{4c^2(a^2 + b^2 - c^2) - (a^2 - b^2)^2} \right] \dots\dots(3).$$

Then area required  $= (8500 \pm 6516.901) \div 2 = 750.84\frac{1}{2}$  or 99.155 acres.

The second is the value required ; the other is for point within the field.



II. Solution by G. B. M. ZERR, A. M., Ph. D., Professor of Mathematics and Applied Science, Texarkana College, Texarkana, Arkansas-Texas.

Let  $ABCD$  be the square,  $OA = 70 = a$ ,  $OB = 40 = c$ ,  $OC = 60 = b$ ,  $O$  the origin,  $(x, y)$  co-ordinates of  $A$ ,  $(u, v)$  co-ordinates of  $C$ ,  $\angle ABE = \theta$ ,  $\angle EBC = \frac{\pi}{2} - \theta$ .

$$\therefore (x - c)^2 + y^2 = (u - c)^2 + v^2, \quad x^2 + y^2 = a^2, \quad u^2 + v^2 = b^2 \dots\dots\dots(1, 2, 3).$$